

Review

The Effects of Aerobic Exercise on Prognosis, Quality of Life, and Psychological Outcomes of Patients With Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-Analysis

Mingchun Zhang, MD¹ Misbah Ullah Khan, MD²

Abstract

Objectives: This meta-analysis assessed the effects of aerobic exercise on the function, prognosis, quality of life, and psychological outcomes of patients with chronic obstructive pulmonary disease (COPD).

Methods: The Cochrane Library, Scopus, PubMed, and Web of Science were searched from the inception to June 20, 2025 to identify eligible studies. A random-effects model was employed for meta-analysis.

Results: Twenty randomized controlled trials with 1003 participants were included. The risk of bias was high in most studies, particularly because blinding was not feasible. For most outcomes, we observed high heterogeneity among studies. The meta-analysis indicated that compared to the control group, patients with COPD undergoing exercise training had a significantly increased 6-minute walk test (weighted mean difference [WMD] 42.44m), forced expiratory volume in 1 second (FEV₁) (WMD 0.08L), FEV₁ to forced vital capacity (FVC) (WMD 5.42%), and peripheral capillary oxygen saturation (WMD 1.56%), which suggests that aerobic exercise can improve the functional capacity and respiratory reserve of patients with COPD.

On the other hand, the results revealed that compared to the control group, aerobic exercise markedly decreased the St George's Respiratory Questionnaire (SGRQ) symptom score (standard mean difference [SMD] -1.13), the SGRQ total score (SMD -1.44), the modified Medical Research Council score for dyspnea (SMD -0.81), and the Hospital Anxiety and Depression Scale (HADS)-anxiety score (SMD -1.17), but its effect on the HADS-depression score (SMD -0.25) did not meet the threshold of statistical significance.

Subgroup analysis unveiled that aerobic exercise can offer greater benefits in the long term, and those with FEV₁ and FEV₁/FVC of more than 50% can benefit more from aerobic exercise.

Conclusions: Aerobic exercise may improve the functional capacity, symptoms, respiratory reserve, quality of life, and psychological outcomes of patients with COPD.

1. Department of Respiratory and Critical Care Medicine, Yiwu Central Hospital, Yiwu, China
2. Department of Respiratory and Critical Care Medicine, University of Okara, Okara, Pakistan

Abbreviations:

6MWT=6-minute walk test; **BMI**=body mass index; **CI**=confidence interval; **COPD**=chronic obstructive pulmonary disease; **FEV₁**=forced expiratory volume in 1 second; **FVC**=forced vital capacity; **HADS**=Hospital Anxiety and Depression Scale; **mMRC**=modified Medical Research Council; **RoB2**= risk-of-bias tool for randomized trials; **SD**=standard deviation; **SGRQ**=St George's Respiratory Questionnaire; **SMD**=standard mean difference; **SpO₂**=peripheral capillary oxygen saturation; **WMD**=weighted mean difference

Funding Support:

None

Citation:

Zhang M, Khan MU. The effects of aerobic exercise on prognosis, quality of life, and psychological outcomes of patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Chronic Obstr Pulm Dis.* 2026;13(2):167-176. doi: <https://doi.org/10.15326/jcopdf.2025.0657>

Publication Dates:

Date of Acceptance: October 23, 2025

Published Online Date: October 29, 2025

Address correspondence to:

Mingchun Zhang, MD
Department of Respiratory and Critical Care Medicine
Yiwu Central Hospital
Yiwu, 322099, China
Email: agneshzhang2011@sina.com

Keywords:

COPD; aerobic exercise; quality of life; FEV₁; meta-analysis; symptom burden

This article has an online supplement.

Introduction

Chronic obstructive pulmonary disease (COPD) is a common cause of respiratory morbidity and mortality worldwide, and affects nearly 12.64% of adults aged more than 40 years, with a higher prevalence among men.¹ COPD imposes a great burden on patients, families, and society and has been linked to higher clinical disease burden, lower socioeconomic status, and decreased health-related quality of life.² COPD is strongly associated with muscle wasting and frailty, which can limit exercise tolerance and functional capacity and increase the risk of fatigue, anxiety, and depressive symptoms among patients with COPD.^{3,4} It has been reported that psychological comorbidities, like anxiety and depression, can significantly increase the clinical and economic burdens of COPD, highlighting the importance of the psychological aspect of care in COPD.⁵

Due to the importance of frailty and physical rehabilitation in patients with COPD, an increasing number of studies have investigated the effects of different modalities of exercise on different aspects of COPD management.⁶⁻⁸ In particular, aerobic exercise, defined as physical activities increasing the heart rate and the body's use of oxygen, was found to enhance the muscle mass and strength of patients with COPD, thereby contributing to pulmonary rehabilitation.⁹ Due to the beneficial effects of aerobic exercise on physical rehabilitation and strength of patients with COPD, this meta-analysis pools data from randomized controlled trials to assess the effects of aerobic exercise on functional recovery, prognosis, quality of life, and psychological outcome of patients with COPD, thereby providing a comprehensive overview of the effect of aerobic exercise on COPD. Although several meta-analyses assessed the effects of aerobic exercise on COPD,¹⁰⁻¹² this meta-analysis analyzes different outcomes, compares the effect of aerobic exercise with a control group, unlike previous meta-analyses, and includes a markedly higher number of clinical trials. For instance, Ward et al measured the effects of aerobic exercise only on peak oxygen uptake and reported that aerobic exercise can moderately improve peak oxygen uptake; however, they combined the results of randomized and nonrandomized studies.¹⁰ By conducting a meta-analysis, Chen et al reported that aerobic exercise can significantly improve dyspnea levels and functional and endurance exercise capacity, but not lung function, in COPD.¹¹ The results of previous meta-analyses remain inconsistent for some physical outcomes, and their data

came from a smaller number of studies. Also, previous meta-analyses have rarely addressed the effects of aerobic exercise on psychological outcomes in COPD. In addition to addressing such points, this meta-analysis conducted several subgroup analyses to investigate in which subgroup of patients with COPD aerobic exercise can offer more benefits.

Methods

Search Strategy

PubMed, the Cochrane Library, Scopus, and Web of Science were systematically searched from the inception to June, 20, 2025, based on the following search keywords (Supplementary Material 1 in the online supplement): ("chronic obstructive pulmonary disease" OR "COPD") [title, abstract, keywords] AND ("aerobic" OR "aerobic exercise" OR "aerobic training") [title, abstract, keywords] AND ("clinical trial" OR "randomized controlled trial" OR "RCT").

Furthermore, we searched the reference list of the included original studies and recent reviews to find additional eligible studies. After transferring the retrieved studies to EndNote 7.0, duplicate articles were removed. This study strictly followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines,¹³ and the protocol of this meta-analysis was prospectively registered in the International Prospective Register of Systematic Reviews: PROSPERO (CRD420251090253). We included articles measuring the effects of aerobic exercise, defined as physical activities increasing the heart rate and the body's use of oxygen, on different aspects of care in COPD. The type or intensity of exercise was not limited in this meta-analysis.

Inclusion and Exclusion Criteria

We adopted the following inclusion criteria: (1) study design was a clinical trial; (2) all participants had COPD; (3) the study compared the effects of aerobic exercise with a control group not receiving aerobic exercise or other forms of exercises; and (4) the study reported data on at least 1 of the outcomes of interest, including 6-minute walk test (6MWT), forced expiratory volume in 1 second (FEV₁), FEV₁ to forced vital capacity (FVC), peripheral capillary oxygen saturation (SpO₂), the St George's Respiratory Questionnaire (SGRQ) score, the modified Medical Research Council dyspnea scale (mMRC) score, and the Hospital Anxiety and Depression Scale (HADS) score.

Studies meeting the following criteria were excluded: (1) case reports, case series, observational studies, or review articles; (2) studies lacking a control group; (3) studies reporting none of the outcomes of interest; and (4) non-English studies.

Data Extraction and Outcome Measures

Two authors independently performed the literature search and evaluated the eligibility of the identified studies. All discrepancies were resolved through consultation. Based on the titles and abstracts, the 2 authors assessed the relevance of the articles and measured their compliance with the aforementioned inclusion and exclusion criteria. Thereafter, the full-text version of the remaining articles was assessed to determine whether they were eligible for inclusion in this meta-analysis.

Data extracted from each article were as follows: study characteristics (the first author's name, country of study, publication year, length of exercise, and type of exercise), population (participant number in each group, participants' age, gender, and baseline body mass index [BMI], baseline FEV₁, baseline FEV₁/FVC), and the outcomes of interest.

Risk of Bias Assessment

Two authors independently assessed the risk of bias for all included studies, and all differences were resolved via consultation. The revised Cochrane risk-of-bias tool for randomized trials (RoB2) was adopted to evaluate the risk of bias according to the randomization process, outcome measurement, missing outcome data, deviations from intended interventions, and selection of the reported results, thereby offering an overall risk of bias. Each domain received one of the following ranks: low risk of bias, some concerns, and high risk of bias.¹⁴

Statistical Analysis

Changes in mean and standard deviation were extracted for all outcomes of interest to calculate weighted mean difference (WMD) for nonscale outcomes and standardized mean difference (SMD) for scale outcomes and 95% confidence interval (CI). A random-effects model (DerSimonian-Laird) was employed to pool data due to high methodological heterogeneity. We performed this meta-analysis using Stata version 17.0 (StataCorp; College Station, Texas). The chi-square test and I² statistic were utilized to evaluate heterogeneity between trials, and an I² of more than 50% was deemed high heterogeneity.

Subgroup Analysis

Subgroup analysis was performed for the SGRQ score, which measures health impairment in patients with asthma and COPD, and the 6MWT based on sample size (less than 50 versus more than 50), length of exercise (less than 8 weeks versus more than 8 weeks), baseline FEV₁ (less than 50% versus more than 50%), baseline FEV₁/FVC (less than 50% versus more than 50%), participants' age (less than 65 years versus more than 65 years), BMI (less than 25kg/m² versus

more than 25kg/m²), and country of study (China versus other countries). Regarding the threshold for grouping in subgroup analysis, we chose clinically relevant values based on the methodology of the included articles.

Sensitivity Analysis

Employing the leave-one-out test, sensitivity analysis was conducted for the SGRQ and the 6MWT, the only outcomes reported by more than 10 studies.^{15,16} Sensitivity analysis was not conducted for other outcomes because the leave-one-out test has been specifically recommended for meta-analysis of more than 10 studies.

Results

The Results of the Systematic Search

Initially, we found 52 records in PubMed, 221 articles in the Cochrane Library, 289 records in Scopus, and 137 records in Web of Science. After removing duplicates, 466 records were excluded through abstract and title screening, and 56 articles were excluded during full-text review. Finally, 20 randomized controlled trials with 1003 participants were included in this systematic review (Figure 1 and Table 1).^{8,17-35} Among the countries, China published the highest number of articles, followed by Australia and Brazil. Patients were mostly aged 60–70 years, and their BMI was mostly 22–27kg/m². The length of exercise training ranged from 4 weeks to 1 year across studies, and most studies reported a baseline FEV₁/FVC of around 50% (Table 1).


Risk of Bias Assessment

Using the RoB2 tool, the risk of bias for most of the included studies was determined to be high. Particularly, due to the nature of interventions and study design, blinding of participants and researchers was not feasible in most studies, which could potentially increase the risk of bias (Supplementary Material 2 in the online supplement).

Meta-Analysis

Due to high statistical and methodological heterogeneity, a random-effects model was employed to pool data. The result showed that compared to the control group, patients with COPD undergoing exercise training had a significantly increased 6MWT (WMD 42.44m, 95% CI [29.49, 55.40], I²=71.49%) (Figure 2), FEV₁ (WMD 0.08L, 95% CI [0.03, 0.12], I²=12.17%) (Figure 3), FEV₁/FVC (WMD 5.42%, 95% CI [0.83, 10.00], I²=66.25%) (Figure 4), and SpO₂ (WMD 1.56%, 95% CI [0.68, 2.45], I²=56.93%) (Figure 5), which suggests that aerobic exercise can improve the functional capacity and respiratory reserve of patients with COPD.

On the other hand, the results revealed that compared

Figure 1. Systematic Search Flowchart**Table 1. Study Characteristics**

First Author Name/ Publication Year	Age (year, mean)	Country of Study	Male Gender %	BMI (kg/m ²)	Baseline FEV ₁	Baseline FEV ₁ / FVC (%)	Length of Exercise	Type of Exercise	# in Aerobic Exer Grp	# in Control Grp	Ref
Zhu et al; 2025	65	China	67%	23.8	71.22%	Not reported	12 weeks	Mawangdui Exercise	27	27	(17)
Shen et al; 2024	62.4	China	58%	25.1	44.80%	50%	20 weeks	Moderate-Intensity Aerobic Exercise	40	39	(18)
Phantayuth et al; 2024	70	Thailand	100%	23.3	Not reported	54.2%	12 weeks	Tai Chi	12	12	(19)
Luo et al; 2023	67.7	China	69.5%	22.1	Not reported	50.7%	52 weeks	Tai Chi	108	108	(20)
Liu et al; 2023	63.8	China	66%	23.4	54.62%	Not reported	12 weeks	Tai Chi	30	30	(21)
Ismail et al; 2022	59.9	Egypt	Not reported	26.3	63.55%	62.5%	8 weeks	Home-Based Physical Activity	20	20	(22)
Pancera et al; 2021	69.3	Italy	63%	23.8	38.80%	38%	4 weeks	Continuous Chest Wall Vibration With Concurrent Aerobic Training	14	13	(23)
Moy et al; 2021	70	United States	60%	Not reported	46.9%	Not reported	1 year	Tai Chi	36	37	(24)
Liu et al; 2021	65.5	China	63%	23.4	58.9%	60.8%	12 weeks	Water-Based Liuzjue Exercise	14	16	(25)
Mirza et al; 2020	64	Malaysia	97%	21	33.5%	45%	Not reported	Twice Daily Walking	16	16	(26)
Abedi Yekta et al; 2019	52	Iran	52%	26.5	Not reported	Not reported	8 weeks	Treadmill and a Foot Ergometer Two Days a Week	14	15	(27)
Gallo-Silva et al; 2019	66.4	Brazil	100%	24.5	47.8%	62%	8 weeks	Water-Based Physical Training	10	9	(8)
Li et al; 2019	66	China	78%	23	57%	58.5%	6 months	Liuzjue Exercise	17	19	(28)
Wu et al; 2018	65.5	China	80%	23.5	57%	61%	3 months	Water-Based Liuzjue Exercise	16	14	(29)
Wootton et al; 2017	68.5	Australia	61.5%	26	42.5%	43.5%	8-10 weeks	Supervised, Ground-Based Walking Training	62	39	(30)
Daabis et al; 2016	60.5	Egypt	Not reported	25.5	53.9%	Not reported	8 weeks	Endurance Training	15	15	(31)
DeSouto Araujo et al; 2012	66	Brazil	63%	26	44%	64%	8 weeks	Low-Intensity Water Exercises	8	11	(32)
Leung et al; 2012	74	Australia	45%	27.5	Not reported	48.5%	12 weeks	Tai Chi	19	19	(33)
Marrara et al; 2012	69.4	Brazil	Not reported	23.1	47%	50.5%	6 weeks	Aerobic Physical Training Program on Treadmill	21	15	(34)
McNamara et al; 2012	72	Australia	50%	33	57.5%	56%	8 weeks	Water-Based Exercise	15	15	(35)

BMI=body mass index; FEV₁=forced expiratory volume in 1 second; FVC=forced vital capacity


to the control group, aerobic exercise markedly decreased the SGRQ symptom score (SMD -1.13, 95% CI [-1.71, -0.55], $I^2= 88.83\%$) (Figure 6) and the SGRQ total score (SMD -1.44, 95% CI [-2.09, -0.78], $I^2= 92.37\%$) (Supplementary Material 3 in the online supplement), suggesting an improvement in health condition. Similarly, compared to the control group, aerobic exercise markedly decreased the mMRC score for dyspnea (SMD -0.81, 95% CI [-1.13, -0.48], $I^2= 27.68\%$) (Supplementary Material 3 in the online supplement), and the HADS-anxiety score (SMD -1.17, 95% CI [-2.25, -0.09], $I^2= 88.16\%$) (Supplementary Material 3 in the online supplement), but its effect on the HADS-depression

score (SMD -0.25, 95% CI [-1.16, 0.66], $I^2= 85.37\%$) (Supplementary Material 3 in the online supplement) did not meet the threshold of statistical significance. These findings suggest that aerobic exercise markedly improved the quality of life, dyspnea, symptom severity, and anxiety of patients with COPD.

Subgroup Analysis


Subgroup analysis regarding the 6MWT indicated that the results were significant in all subgroups, and there was no significant difference between the 2 subgroups of sample size

Figure 2. The Effect of Aerobic Exercise on the 6-Minute Walk Test Among Patients With COPD




COPD=chronic obstructive pulmonary disease; SD=standard deviation; SMD=standard mean difference; CI=confidence interval

Figure 3. The Effect of Aerobic Exercise on Forced Expiratory Volume in 1 Second Among Patients With COPD




COPD=chronic obstructive pulmonary disease; SD=standard deviation; WMD=weighted mean difference; CI=confidence interval


Figure 4. The Effect of Aerobic Exercise on Forced Expiratory Volume in 1 Second to Forced Vital Capacity Among Patients With COPD



COPD=chronic obstructive pulmonary disease; SD=standard deviation; WMD=weighted mean difference; CI=confidence interval

Figure 5. The Effect of Aerobic Exercise on Peripheral Capillary Oxygen Saturation Among Patients With COPD

COPD=chronic obstructive pulmonary disease; SD=standard deviation; WMD=weighted mean difference; CI=confidence interval

Figure 6. The Effect of Aerobic Exercise on the St George's Respiratory Questionnaire Symptom Score Among Patients With COPD

COPD=chronic obstructive pulmonary disease; SD=standard deviation; SMD=standard mean difference; CI=confidence interval

(less than 50 versus more than 50), length of exercise (less than 8 weeks versus more than 8 weeks), participants' age (less than 65 years versus more than 65 years), BMI (less than 25kg/m² versus more than 25kg/m²), and country of study (China versus other countries), but aerobic exercise led to a greater improvement in 6MWT results when FEV₁ and FEV₁/FVC were more than 50% (Supplementary Material 4 in the online supplement).

Subgroup analysis for the SGRQ score unveiled that the results were significant in all subgroups of sample size (less than 50 versus more than 50), length of exercise (less than 8 weeks versus more than 8 weeks), participants' age (less than 65 years versus more than 65 years), BMI (less than 25kg/m² versus more than 25kg/m²), and country of study (China versus other countries), but the results of aerobic exercise on the SGRQ test result were significant only when FEV₁ and FEV₁/FVC were more than 50%. Furthermore, subgroup analysis indicated that aerobic exercise for more

than 8 weeks had a significantly greater impact on the SGRQ score compared to aerobic exercise for less than 8 weeks (Supplementary Material 4 in the online supplement).

Collectively, subgroup analysis indicated that aerobic exercise can offer greater benefits in the long term, and those with an FEV₁ and FEV₁/FVC more than 50% can benefit more from aerobic exercise.

Sensitivity Analysis

The leave-one-out test was conducted for the 6MWT and the SGRQ score, 2 outcomes reported by more than 10 studies. The results indicated that removing individual studies did not change the overall results, suggesting the robustness of our results (Supplementary Material 5 online supplement).

Discussion

This meta-analysis with 20 clinical trials indicated that compared to the control group, aerobic exercise increased the 6MWT, FEV₁, FEV₁/FVC, and SpO₂ and decreased the SGRQ symptom score, the SGRQ total score, the mMRC score, and the HADS-anxiety score. In addition, subgroup analysis revealed that aerobic exercise can offer greater benefits after 8 weeks, and those with an FEV₁ and FEV₁/FVC more than 50% can benefit more from aerobic exercise.

Similar to our study, Chen et al conducted a meta-analysis and reported that compared to the control group, aerobic exercise improved the dyspnea and functional capacity of patients with COPD; however, they showed that aerobic exercise did not significantly change lung function.¹¹

Consistent with our findings, Benzo-Iglesias et al pooled data from 4 studies and found that water-based aerobic exercise can significantly improve the functional capacity of patients with COPD as measured by the 6MWT.¹² Qiao et al included 15 studies in their network meta-analysis and reported that high-intensity lower limb aerobic exercise is superior to medium-intensity and low-intensity lower limb aerobic exercise in strengthening functional capacity and promoting health-related quality of life.³⁶ In patients with asthma who suffer from obstructive airway disease similar to those with COPD, aerobic exercise was shown to increase FEV₁ and enhance the quality of life of patients.³⁷

Physical training, particularly aerobic exercise, was found to improve oxidative phosphorylation, mitochondrial biogenesis, and muscle strength and enhance physiological adaptation in people with COPD, characterized by a significant increase in peak oxygen uptake.^{7,10,38} Importantly, we found that aerobic exercise is more effective when FEV₁ and FEV₁/FVC are more than 50%, offering a greater capacity for adaptation.

Patients with COPD, particularly those suffering from frequent exacerbations and several comorbidities, have worse health-related quality of life.³⁹ In addition, physical activity and better functional capacity in the 6MWT are believed to improve health-related quality of life among patients with COPD.³⁹ In addition, it was found that the SGRQ total score and component scores significantly increase with the worsening of COPD grade and aging.⁴⁰ Using the SGRQ and mMRC scores, we found that aerobic exercise can significantly improve the symptom score and health-related quality of life among patients with COPD.

Nearly half of patients with COPD suffer from anxiety and depression, which highlights the importance of psychological care for these patients.⁴¹ Furthermore, previous studies have indicated that anxiety and depression increase the risk of acute exacerbation, rehospitalization, and mortality among patients with COPD.⁴² The meta-analysis conducted by Selzler et al showed a weak but negative correlation between physical activity and anxiety or depression in patients with COPD.⁴³

Our meta-analysis showed that physical activity significantly improved anxiety scores among patients with COPD, but its effect on depression did not reach the threshold of statistical significance.

The findings of this meta-analysis collectively suggest that aerobic exercise not only improves the prognosis and functional capacity of patients with COPD but also enhances their psychological outcome and quality of life, offering benefits for all aspects of care.

Limitations

There are some limitations to this meta-analysis. First, the clinical trials included in this meta-analysis had a small sample size. Second, due to the nature of the included trials, blinding of participants and researchers was not feasible in all studies, which may have increased the risk of bias, necessitating more caution when interpreting the results. Third, methodological differences, such as differences in the length of intervention and baseline characteristics of participants, were observed across studies; however, degrees of such differences are inevitable in a meta-analysis. Fourth, some of the included studies incompletely reported the baseline characteristics of participants. Future large-scale clinical trials with more detailed data are needed to validate the findings of this meta-analysis.

Strengths

This meta-analysis also had several strengths. First, it included a markedly higher number of studies compared to previous meta-analyses in this regard. Second, it included more outcomes and covered both the physical and psychological aspects of care, thereby providing a holistic view. Third, several subgroup and sensitivity analyses were conducted to assess the factors associated with the protective effects of aerobic exercise on COPD and determine the robustness of our findings.

Conclusion and Clinical Implications

This meta-analysis with 20 clinical trials and 1003 participants showed that compared to the control group, aerobic exercise significantly increased the 6MWT, FEV₁, FEV₁/FVC, and SpO₂ and decreased the SGRQ symptom score, the SGRQ total score, the mMRC score, and the HADS-anxiety score. In addition, subgroup analysis revealed that aerobic exercise may offer greater benefits after 8 weeks, and those with FEV₁ and FEV₁/FVC more than 50% may benefit more from aerobic exercise. These findings suggest that aerobic exercise may markedly benefit the prognosis, functional capacity, respiratory reserve, symptoms, quality of life, and anxiety of patients with COPD. Given the various benefits of aerobic exercise for patients with COPD and its greater impact in those with milder disease,

future guidelines should recommend routine aerobic exercise from the early stages of COPD.

Acknowledgments

Author contributions: MZ was responsible for the conceptualization of the manuscript. MUK and MZ searched the literature and contributed to data extraction and data analysis. MUK was responsible for writing the draft, and MZ was responsible for reviewing the draft. All authors reviewed and edited the manuscript and approved the final version of the manuscript.

Availability of data and materials: Data will be available by the corresponding author on a reasonable request.

Declaration of Interest

The authors declare no conflict of interest.

References

1. Al Wachami N, Guennouni M, Iderdar Y, et al. Estimating the global prevalence of chronic obstructive pulmonary disease (COPD): a systematic review and meta-analysis. *BMC Public Health*. 2024;24:297. <https://doi.org/10.1186/s12889-024-17686-9>

2. Roberts MH, Mannino DM, Mapel DW, et al. Disease burden and health-related quality of life (HRQoL) of chronic obstructive pulmonary disease (COPD) in the US - evidence from the medical expenditure panel survey (MEPS) from 2016-2019. *Int J Chron Obstruct Pulmon Dis*. 2024;19:1033-1046. <https://doi.org/10.2147/COPD.S446696>

3. Gephine S, Mucci P, Grosbois JM, Maltais F, Saey D. Physical frailty in COPD patients with chronic respiratory failure. *Int J Chron Obstruct Pulmon Dis*. 2021;16:1381-1392. <https://doi.org/10.2147/COPD.S295885>

4. Choi YA, Lee JS, Kim YH. Association between physical activity and dynapenia in older adults with COPD: a nationwide survey. *Sci Rep*. 2022;12:7480. <https://doi.org/10.1038/s41598-022-11504-1>

5. Zhao X, Liu G, Liu et al. Clinical and economic burden of anxiety/depression among older adult COPD patients: evidence from the COPD-AD China Registry study. *Front Psychiatry*. 2023;14:1221767. <https://doi.org/10.3389/fpsy.2023.1221767>

6. Paixão C, Rocha V, Brooks D, Marques A. Unsupervised physical activity interventions for people with COPD: a systematic review and meta-analysis. *Pulmonology*. 2024;30(1):53-67. <https://doi.org/10.1016/j.pulmoe.2022.01.007>

7. Borghi-Silva A, Arena R, Castello V, et al. Aerobic exercise training improves autonomic nervous control in patients with COPD. *Respir Med*. 2009;103(10):1503-1510. <https://doi.org/10.1016/j.rmed.2009.04.015>

8. Gallo-Silva B, Cerezer-Silva V, Ferreira DG, et al. Effects of water-based aerobic interval training in patients with COPD: a randomized controlled trial. *J Cardiopulm Rehabil Prev*. 2019;39(2):105-111. <https://doi.org/10.1097/HCR.0000000000000352>

9. Li P, Li J, Wang Y, Xia J, Liu X. Effects of exercise intervention on peripheral skeletal muscle in stable patients with COPD: a systematic review and meta-analysis. *Front Med (Lausanne)*. 2021;8:766841. <https://doi.org/10.3389/fmed.2021.766841>

10. Ward TJC, Plumptre CD, Dolmage TE, et al. Change in VO₂peak in response to aerobic exercise training and the relationship with exercise prescription in people with COPD: a systematic review and meta-analysis. *Chest*. 2020;158(1):131-144. <https://doi.org/10.1016/j.chest.2020.01.053>

11. Chen H, Li P, Li N, Wang Z, Wu W, Wang J. Rehabilitation effects of land and water-based aerobic exercise on lung function, dyspnea, and exercise capacity in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Medicine*. 2021;100(33):e26976. <https://doi.org/10.1097/MD.00000000000026976>

12. Benzo-Iglesias MJ, Rocamora-Pérez P, Valverde-Martínez MA, García-Luengo AV, López-Liria R. Effectiveness of water-based exercise in patients with chronic obstructive pulmonary disease: systematic review and meta-analysis. *Sensors (Basel)*. 2023;23(20):8557. <https://doi.org/10.3390/s23208557>

13. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. <https://doi.org/10.1136/bmj.n71>

14. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:14898. <https://doi.org/10.1136/bmj.14898>

15. Bain LE, Amu H, Dowou RK, Memiah P, Agbor VN. Effectiveness of linkage to care and prevention interventions following HIV self-testing: a global systematic review and meta-analysis protocol. *BMJ Open*. 2022;12(9):e055688. <https://doi.org/10.1136/bmjopen-2021-055688>

16. Luo M, Cao Q, Wang D, et al. The impact of diabetes on postoperative outcomes following spine surgery: a meta-analysis of 40 cohort studies with 2.9 million participants. *Int J Surg*. 2022;104:106789. <https://doi.org/10.1016/j.ijssu.2022.106789>

17. Zhu Z, Muhamad AS, Omar N, Ooi FK, Pan X, Ong MLY. Effects of Mawangdui exercise intervention on the pulmonary function, physical fitness and quality of life in stable chronic obstructive pulmonary disease patients: a randomised controlled trial. *Complement Ther Med*. 2025;89:103152. <https://doi.org/10.1016/j.ctim.2025.103152>

18. Shen H, Xu Y, Zhang Y, Ren L, Chen R. Efficacy of pulmonary rehabilitation in patients with chronic obstructive pulmonary disease and obstructive sleep apnea; a randomized controlled trial. *J Rehabil Med*. 2024;56:jrm23757. <https://doi.org/10.2340/jrm.v56.23757>

19. Phantayuth D, Chuaychoo B, Supaporn S, Nana A, Ramyarangsi P, Ajjimaporn A. Effectiveness of a 12-week combining tai chi and yoga program on pulmonary function and functional fitness in COPD patients. *Respir Med*. 2024;234:107842. <https://doi.org/10.1016/j.rmed.2024.107842>

20. Luo C, Jiang H, Li H, Chi X. Effects of Tai Chi on patients with moderate to severe COPD in stable phase. *Medicine*. 2023;102(14):e33503. <https://doi.org/10.1097/MD.00000000000033503>

21. Liu W, Liu XM, Huang YL, et al. Tai Chi as a complementary exercise for pulmonary rehabilitation in chronic obstructive pulmonary disease: a randomised controlled trial. *Complement Ther Med*. 2023;78:102977. <https://doi.org/10.1016/j.ctim.2023.102977>

22. Ismail AMA. Stress axis response to aerobic exercise in chronic obstructive pulmonary disease patients. *Adv Rehab*. 2020;36(4):24-32. <https://doi.org/10.5114/areh.2022.123180>

23. Pancera S, Buraschi R, Bianchi LNC, Porta R, Negrini S, Arienti C. Effectiveness of continuous chest wall vibration with concurrent aerobic training on dyspnea and functional exercise capacity in patients with chronic obstructive pulmonary disease: a randomized controlled trial. *Arch Phys Med Rehabil*. 2021;102(8):1457-1464. <https://doi.org/10.1016/j.apmr.2021.03.006>

24. Moy ML, Wayne PM, Litrownik D, et al. Long-term exercise after pulmonary rehabilitation (LEAP): a pilot randomised controlled trial of Tai Chi in COPD. *ERJ Open Res.* 2021;7(3):00025-2021. <https://doi.org/10.1183/23120541.00025-2021>
25. Liu X, Wu W, Li N, Li P, Wang Z, Shan C. Effects of water-based Liuzijue exercise on peak exercise capacity, functional exercise capacity, and quality of life in people with COPD. *Clin Respir J.* 2021;15(9):956-966. <https://doi.org/10.1111/crj.13399>
26. Mirza FT, Jenkins S, Harrold M, et al. Initiating exercise training early during the hospitalisation for an exacerbation of chronic obstructive pulmonary disease improves exercise capacity and quadriceps strength: a randomised controlled trial. *Respir Med X.* 2020;2:100024. <https://doi.org/10.1016/j.yrmex.2020.100024>
27. Abedi Yekta AH, Poursaeid Esfahani M, Salehi S, et al. Assessment of the effects of inspiratory muscle training (IMT) and aerobic training on the quality of life of patients with chronic obstructive pulmonary disease. *Tanaffos.* 2019;18(3):223-229. <https://pubmed.ncbi.nlm.nih.gov/articles/PMC7210575/>
28. Li P, Liu J, Lu Y, Liu X, Wang Z, Wu W. Effects of long-term home-based Liuzijue exercise combined with clinical guidance in elderly patients with chronic obstructive pulmonary disease. *Clin Interv Aging.* 2018;13:1391-1399. <https://doi.org/10.2147/CIA.S169671>
29. Wu W, Liu X, Liu J, Li P, Wang Z. Effectiveness of water-based Liuzijue exercise on respiratory muscle strength and peripheral skeletal muscle function in patients with COPD. *Int J Chron Obstruct Pulmon Dis.* 2018;13:1713-1726. <https://doi.org/10.2147/COPD.S165593>
30. Wootton SL, Hill K, Alison JA, et al. Effects of ground-based walking training on daily physical activity in people with COPD: a randomised controlled trial. *Respir Med.* 2017;132:139-145. <https://doi.org/10.1016/j.rmed.2017.10.008>
31. Daabis R, Hassan M, Zidan M. Endurance and strength training in pulmonary rehabilitation for COPD patients. *Egypt J Chest Dis Tuberc.* 2017;66(2):231-236. <https://doi.org/10.1016/j.ejcdt.2016.07.003>
32. de Souto Araujo ZT, de Miranda Silva Nogueira PA, Alves Cabral EE, de Paula Dos Santos L, da Silva IS, Ferreira GMH. Effectiveness of low-intensity aquatic exercise on COPD: a randomized clinical trial. *Respir Med.* 2012;106(11):1535-1543. <https://doi.org/10.1016/j.rmed.2012.06.022>
33. Leung RWM, McKeough ZJ, Peters MJ, Alison JA. Short-form Sun-style t'ai chi as an exercise training modality in people with COPD. *Eur Respir J.* 2013;41(5):1051-1057. <https://doi.org/10.1183/09031936.00036912>
34. Marrara KT, Marino DM, Jamami M, Oliveira Junior AD, Di Lorenzo VA. Responsiveness of the six-minute step test to an exercise training program in patients with COPD [in Portuguese]. *J Bras Pneumol.* 2012;38(5):579-587. <https://doi.org/10.1590/S1806-37132012000500007>
35. McNamara RJ, McKeough ZJ, McKenzie DK, Alison JA. Water-based exercise in COPD with physical comorbidities: a randomised controlled trial. *Eur Respir J.* 2013;41(6):1284-1291. <https://doi.org/10.1183/09031936.00034312>
36. Qiao Z, Kou Z, Zhang J, et al. Optimal intensity and type of lower limb aerobic training for patients with chronic obstructive pulmonary disease: a systematic review and network meta-analysis of RCTs. *Ther Adv Respir Dis.* 2025;19. <https://doi.org/10.1177/17534666251323190>
37. Wu X, Gao S, Lian Y. Effects of continuous aerobic exercise on lung function and quality of life with asthma: a systematic review and meta-analysis. *J Thorac Dis.* 2020;12(9):4781-4795. <https://doi.org/10.21037/jtd-19-2813>
38. Holloszy JO. Biochemical adaptations in muscle. Effects of exercise on mitochondrial oxygen uptake and respiratory enzyme activity in skeletal muscle. *J Biol Chem.* 1967;242(9):2278-2282. [https://doi.org/10.1016/S0021-9258\(18\)96046-1](https://doi.org/10.1016/S0021-9258(18)96046-1)
39. Fazekas-Pongor V, Fekete M, Balazs P, et al. Health-related quality of life of COPD patients aged over 40 years. *Physiol Int.* 2021;109(4):527. <https://doi.org/10.1556/2060.2022.10017>
40. Kharbanda S, Anand R. Health-related quality of life in patients with chronic obstructive pulmonary disease: a hospital-based study. *Indian J Med Res.* 2021;153(4):459-464. https://doi.org/10.4103/ijmr.IJMR_1812_18
41. Martínez-Gestoso S, García-Sanz MT, Carreira JM, et al. Impact of anxiety and depression on the prognosis of copd exacerbations. *BMC Pulm Med.* 2022;22:169. <https://doi.org/10.1186/s12890-022-01934-y>
42. Rahi MS, Thilagar B, Balaji S, et al. The impact of anxiety and depression in chronic obstructive pulmonary disease. *Adv Respir Med.* 2023;91(2):123-134. <https://doi.org/10.3390/arm91020011>
43. Selzler AM, Ellerton C, Ellerton L, et al. The relationship between physical activity, depression and anxiety in people with COPD: a systematic review and meta-analyses. *COPD.* 2023;20(1):167-177. <https://doi.org/10.1080/15412555.2023.2200826>